

Digestibility of Starter Feeds in Calves: Modeling the effects of liquid intake and weaning on digestibility of nutrients in pre- and post-weaned dairy calves

J. D. Quigley¹, T. M. Hill, F. X. Suarez-Mena, T. S. Dennis, J. M. Aldrich, and R. L. Schlotterbeck

Nurture Research Center, Provimi North America, Cargill Premix and Nutrition

¹ Corresponding author. Email: jquigley@provimi-na.com

■ Take Home Messages

- Digestion of nutrients (particularly carbohydrates) from calf starters depends on rumen development.
- Consumption of more than approximately 700 g/d of milk replacer solids may depress starter intake and rumen development.
- Delayed rumen development reduces post-weaning digestion of nutrients and may predispose calves to reduced growth, increased stress, and potentially increased risk of disease.

■ Introduction

Accurate predictions of nutrient supply and nutrient requirements are essential to modern ration formulations and animal production. Accurate and precise models allow the provision of nutrients to meet requirements for maintenance and optimal production without supplying excess nutrients that contribute to inefficiency or environmental damage.

Most nutrient models predict supply of metabolizable energy (ME) and metabolizable protein (MP); in lactation models, flow of nutrients are predicted from endogenous, microbial, and undegraded dietary sources. Nutrient requirements are usually predicted using factorial calculation of requirements for maintenance (adjusted for environmental and management considerations), growth, pregnancy, and lactation. Only maintenance and growth predictions are used to predict nutrient requirements for calves, with requirements for pregnancy included for primiparous heifers.

For young calves and heifers, prediction of nutrient supply by the 2001 Nutrient Requirements of Dairy Cattle (NRC, 2001) assumes fixed digestibility and metabolizability of energy and protein. For example, calculation of ME from milk replacer is assumed to be the caloric content of protein, fat, and lactose adjusted for digestibility and metabolizability:

$$\text{ME (Mcal/kg)} = [(0.057 \times \text{CP}) + (0.092 \times \text{EE}) + (0.0395 \times \text{CHO})] \times 97\% \times 96\%,$$

where: CP = crude protein %, EE = ether extract %, CHO = carbohydrate %, 97% = digestibility of nutrients, and 96% = metabolizability of digested nutrients.

Metabolizable energy content of calf starters is calculated as the sum of the digestible fractions of protein, non-fibre carbohydrates, neutral detergent fibre (NDF), crude protein (CP), and fat as described in the 2001 Dairy NRC (NRC, 2001) for adult cattle. Neither liquid nor starter feeds are corrected for differences in digestibility caused by age or development of the gastrointestinal tract in these models.

In young calves, digestibility of dry feeds (concentrates and forages) depend on the development of ruminal fermentation and intestinal digestion. This is particularly true for NDF (primarily fermented in the rumen) and starch (dependent on ruminal fermentation and small intestinal digestion). Studies have shown that fibre fermentation is limited in neonatal calves (Chapman et al., 2016; Hill et al., 2016a, b). Further, pancreatic α -amylase production is low at birth (Siddons, 1968) but increases with age (Huber et al., 1961; Morrill et al., 1970) along with total pancreatic secretion (McCormick and Stewart, 1966) thereby affecting small intestinal digestion of starch (Morrill et al., 1970).

Development of microbial fermentation changes the flow of nutrients from the stomach. Prior to weaning, nutrients are derived primarily from milk protein, fat, and lactose; after weaning, nutrients are provided by volatile fatty acids absorbed from the rumen and microbial protein that increases in flow with increasing dry feed intake (Leibholz, 1975; Quigley et al., 1985).

Changing the amounts and types of liquid fed to calves may alter age at which dry feed intake begins (Hill et al., 2006a, b; Strzetelski et al., 2001) thereby altering rumen development. This is particularly true when large amounts of liquid are fed (i.e., greater than about 700 g of solids from liquid/day for Holstein calves) since large amounts of liquid consumed will delay rumen development (Terré, et. al, 2007). Several studies have reported increased BW at weaning for calves fed large amounts of liquid pre-weaning; however, the advantage in growth compared to conventional feeding methods (500-700 g of solids/day) may be lost as BW gain slows dramatically in the period immediately post-weaning. We have attempted to quantify the effects

of increased milk replacer allowance on digestibility of starter and its effects on growth and efficiency of young calves to determine if differences in the digestion of nutrients, particularly of carbohydrates, might be at least partially responsible for differences in growth.

■ Digestion of Solid Feed

Calves are commonly weaned between 1 and 3 months of age in most dairy systems, with the most common age being approximately 9 weeks of age in the U.S. (USDA, 2016). Weaning to dry feed requires that the calf has sufficient digestive and fermentative capability to provide nutrients to support maintenance and growth. Further, the source of nutrients changes from milk digested primarily in the small intestine to grain-based ingredients fermented in the rumen and (or) digested in the small intestine. Therefore, gastrointestinal, hepatic, and systemic enzyme systems must be sufficiently adapted to changing sources of nutrients. If a calf is inadequately prepared for weaning, performance may suffer and predispose calves to reduced growth, poor efficiency, and even increased susceptibility to disease (Roth et al., 2008, 2009).

The most important factor in promoting rumen development and adaptation in preparation for weaning is consumption of dry feed containing fermentable carbohydrates – particularly sugars and starches – that are fermented to propionate and butyrate in the rumen by resident rumen bacteria. Production of volatile fatty acids and microbial protein stimulate a series of adaptations in the rumen, gastrointestinal tract, hepatic tissues, and systemically that promote gluconeogenesis, production and release of β -hydroxybutyrate by rumen epithelium, and utilization of acetate by peripheral tissues (Howarth et al., 1968; Huber, 1969; Baldwin et al., 2004).

In the past 15 years, some dairy experts have recommended feeding milk or milk replacer in excess of the traditional recommendations (approximately 10% of body weight as milk or reconstituted milk replacer) to increase rate of gain and take advantage of improved calf efficiency (Diaz et al., 2001; Davis-Rincker et al., 2011; Moallem et al., 2010). High digestibility and metabolizability of liquid feeds compared to higher fibre ingredients in calf starters naturally contribute to greater efficiency of BW gain.

Calves fed whole milk ad libitum or milk replacer to amounts >1 kg of powder per day gain impressive amounts of BW. For example, Jasper and Weary (2002) reported that calves fed milk ad libitum were 8 kg heavier at the end of a 63-d feeding period compared to calves fed milk at 10% of BW. All calves were weaned at 42 d. However, daily BW gains in calves fed ad libitum were markedly lower during the week of weaning (0.36 vs. 0.53 kg) and after weaning (0.68 vs. 0.85 kg) so that BW differences at 63 d were not as great as the difference prior to weaning.

Differences in growth rate post-weaning in calves fed differently pre-weaning may be due to differences in gastrointestinal development and digestion. Several recent studies indicate that digestion of nutrients from dry feeds varies when calves are fed varying amounts of liquid pre-weaning. Terré et al. (2007) fed Holstein bull calves (19 d of age at start of the trial) milk replacer (MR) at levels typical of conventional feeding (CF; 4 L/d with weaning at 35 d of the study) or an enhanced feeding (EF) program wherein the amount of MR was increased to 7 L/d and then reduced to weaning. Total starter intake on the CF and EF programs prior to weaning were 23.8 and 12.6 kg, respectively. Results of a digestion trial conducted during d 38-42 of the study are in Table 1. These data indicate clearly that digestion of dry feed was impaired in calves fed EF, likely due to inadequate rumen development as a result of lower starter intake.

Table 1. Apparent total tract digestibility of dry feed in calves fed 4 L/d of milk replacer (MR) at 12.5% DM dilution rate from d 1–28, and 2 L/d from d 29 to d 35 (CF) or MR at 18% DM dilution rate: 4 L/d from d 1–6, 6 L/d from d 7–13, 7 L/d from d 14–20, 6 L/d from d 21–28, and 3 L/d from d 29 to 35 (EF). Digestibility was measured the week after weaning. Adapted from Terré et al. (2007).

Digestibility, %	CF	EF	SE	P
Dry matter	77.4	71.8	1.23	0.01
Organic matter	78.7	73.2	1.18	0.01
Crude protein	77.1	71.6	1.29	0.01
Neutral detergent fibre	34.7	20.3	3.79	0.02
Gross energy	75.6	69.8	1.25	0.01

Digestion of NDF (derived primarily from wheat middlings, soybean hulls, and wheat distiller's grains) in the study by Terré et al. (2007) was lower in EF calves compared to CF calves (20.3 vs. 34.7%; Table 1). Since the disappearance of NDF is due primarily to ruminal fermentation, it is likely that reduced NDF digestion was due to inadequate or incomplete rumen fermentation in EF calves. Reduced NDF digestibility occurred in EF calves in spite of a higher rumen pH (5.73 vs. 5.99). Rumen pH less than approximately 6.0 is associated with impaired ruminal fibre fermentation (Allen, 1997; Shriver et al., 1986) due to the pH sensitivity of cellulolytic bacteria in the rumen (Hoover, 1986; Russell et al., 1996). In the study by Terré et al. (2007), the authors attributed higher ruminal pH to lower ruminal activity due to lower starter intake and a lack of substrate available for fermentation.

Leibholz (1975) monitored digestion of nutrients in calves fed whole milk or MR to weaning at 35 d of age. After weaning, calves were offered a pelleted feed consisting of 58% barley, 20% soybean meal, 15% wheat straw, 3%

molasses plus vitamins and minerals. The diet contained 15% protein and 13% ADF; we estimated the diet contained 2.7 Mcal of ME/kg and 50% non-fibre carbohydrate.

By 6 wk of age (1 wk post-weaning), digestibility of ADF reached 57% and did not change markedly thereafter. However, the site of ADF digestion changed dramatically with time after weaning as most ADF was digested in the hindgut during the first 4 wk of the trial (Figure 1).




Figure 1. Digestion of acid detergent fiber in calves fed milk or milk replacer to weaning at 5 wk of age. Digestion was measured in the stomach and intestines using duodenally cannulated calves. Adapted from Leibholz, 1975.

Weekly DMI for each week of the 8 wk study were 0.6, 1.1, 1.5, 2.1, 2.2, 2.4, 2.5 and 2.5 kg/d. Intake of ADF ranged from 77 g/d in the 1st week post-weaning to 325 g/d at wk 8. Therefore, it is possible that the higher digestion of ADF in the hindgut during the first few weeks after weaning was due to small amounts of ADF consumed.

Hill et al. (2010) fed calves (2-3 d of age at start of study) one of four MR programs: 0.44 kg of DM of a 21% CP, 21% fat MR powder fed daily for 42 d (A); 0.66 kg of DM of a 27% CP, 17% fat MR powder fed daily for 42 d (B); 0.66 kg of DM of a 27% CP, 17% fat MR powder daily fed for 28 d (C); or up to 1.09 kg of DM of a 29% CP, 21% fat MR daily fed for 49 d (D). Digestibility

estimates were made on d 53 to 56. Table 2 shows clearly that digestion of dry matter (DM) and organic matter (OM) were lower when calves were fed large amounts of MR prior to weaning (treatment D). During the digestibility period (d 53 to 56), intake of starter DM was 2.2, 2.3, 2.5 and 1.9 kg/d for treatments A, B, C, and D, respectively. The trend ($P < 0.08$) for low starter DM intake coupled with significantly lower digestion of DM resulted in calves on treatment D only consuming about 71% of the digestible DM of calves on the other treatments.

Table 2. Total tract apparent digestion of dry matter (DM), organic matter (OM), crude protein (CP) or fat in calves fed one of four MR programs: 0.44 kg of DM of a 21% CP, 21% fat MR powder fed daily for 42 d (A); 0.66 kg of DM of a 27% CP, 17% fat MR powder fed daily for 42 d (B); 0.66 kg of DM of a 27% CP, 17% fat MR powder daily fed for 28 d (C); or up to 1.09 kg of DM of a 29% CP, 21% fat MR daily fed for 49 d (D). Adapted from Hill et al., 2010.

Digestion, %	A	B	C	D	SE	P
DM	75.6 ^a	78.3 ^a	78.7 ^a	67.3 ^b	2.19	0.01
OM	77.4 ^a	78.3 ^a	78.7 ^a	68.0 ^b	2.20	0.01
CP	72.4	72.3	74.1	71.8	2.58	0.83
Fat	70.3	75.4	76.3	75.4	3.37	0.33

^{a,b}Means in the same row with different superscripts differ, $P < 0.05$.

More recently, Chapman et al. (2016) reported that digestion of nutrients, particularly of NDF and ADF, were reduced during the digestion period of d 52-58 of age when calves were fed MR up to 0.87 kg/d (Table 3). Although digestion of all nutrients (except starch) were reduced significantly, digestion of NDF and ADF were reduced nearly 50% in calves fed large amounts of milk pre-weaning.

Conversely, Chapman et al. (2017) reported no difference in NDF digestion when calves were fed milk replacer at 446, 669, or 892 g/d of milk replacer during the digestibility measurement period. Further, NDF digestion was 58, 69, and 69%, respectively, suggesting extensive digestion of fibre by the calves. However, the starter used in the study contained only 16% NDF and starter intake during the trial was 1.1, 0.7 and 0.4 kg/d, respectively. Measurements were taken prior to weaning, which may have increased the error associated with measurement.

Table 3. Body weight (BW), DM intake (DMI) and total tract digestibility of nutrients in calves fed conventional [CON; 0.44 kg of dry matter (DM) 21% crude protein (CP), 21% fat powder fed for 42 d], moderate (MOD; 0.66 kg of DM 27% CP, 17% fat powder fed for 42 d), and aggressive program (AGG; up to 0.87 kg of DM 27% CP, 17% fat powder fed for 49 d). Digestibility was measured from d 51-56. From Chapman et al., 2016.

Item	CON	MOD	AGG	SE	P
BW, kg	62.7 ^a	72.3 ^b	82.8 ^c	4.05	0.01
DMI, kg/d	2.04	2.30	2.28	0.258	0.08
Digestibility, %					
DM	77.6 ^a	76.9 ^a	66.0 ^b	1.67	0.01
OM	79.2 ^a	78.2 ^a	67.9 ^b	1.65	0.01
ADF	56.3 ^a	53.2 ^a	26.7 ^b	3.89	0.01
NDF	54.1 ^a	50.7 ^a	26.2 ^b	2.86	0.01
Starch	96.7	94.5	94.0	1.33	0.36
CP	71.9 ^a	74.1 ^a	56.3 ^b	2.72	0.02
Sugar	93.1 ^a	91.5 ^a	86.2 ^b	1.68	0.02
Fat	81.4 ^a	83.2 ^a	74.1 ^b	1.84	0.01

^{a,b,c}Means in the same row with different superscripts differ, $P < 0.05$.

Finally, Dennis et al. (2018) compared elevated MR feeding programs and methods of weaning on temporal changes in nutrient digestibility in pre- and post-weaned calves. In this study, calves were fed 0.66 kg of DM from MR from d 0 to 39, 0.33 kg from d 40 to 42 (MOD); 0.87 kg from d 0 to 4, 1.09 kg from d 5 to 35, 0.54 kg from d 36 to 42 (HIGH6); 0.87 kg from d 0 to 4, 1.09 kg from d 5 to 45, 0.54 kg from d 46 to 53 (HIGH8); or 0.87 kg for 4 d, 1.09 kg from d 5 to 39, 0.87 kg from d 40 to 43, 0.66 kg from d 44 to 47, 0.44 kg from d 48 to 50, 0.22 kg from d 51 to 53 (GRAD8). Calves were fed starter (21% CP, 37% starch, 3.7% fat) to 56-d, then 95% starter plus 5% hay (7% CP, 64% NDF) to d 112. Digestion of NDF and ADF were consistently higher in calves fed MOD6 compared to other treatments to 84 d. Only at 112 d of age were digestibility coefficients similar among treatments, suggesting that level of MR feeding can depress nutrient digestion from dry feed post-weaning and that weaning strategy (6 vs. 8 wk of age and abrupt vs. gradual) had little effect on this depression.

The majority of these data suggest that calves fed large amounts of milk pre-weaning may have difficulty digesting nutrients from dry feed during the immediate post-weaning period. There are numerous implications to these findings. For example, digestion of starters containing greater amounts of

fibrous by-products may be difficult if calves are fed large amounts of liquid pre-weaning. Also, it may be necessary to use increasingly complex liquid reduction strategies to ensure that starter intake (and digestibility) is adequate prior to weaning.

Because fibre digestion is primarily influenced by cellulolytic fermentation in the rumen, the low digestibility of ADF and NDF (Table 3) indicate that the rumen is less well developed in calves fed greater amounts of MR (Chapman et al., 2016). Also, fibre digesting microorganisms are established in the rumen more slowly than starch and sugar digesting microorganisms (Anderson et al., 1987). Finally, selection of ingredients that may negatively affect rumen fermentation (e.g., inclusion of oil-containing ingredients) may also reduce total DM digestion (Hill et al., 2015).

To better understand the changes in NDF digestion with age and diet, Hill et al. (2016b) fed calves a moderate or aggressive milk replacer feeding program and monitored changes in nutrient digestion with advancing age. Figure 2 shows changes in NDF digestion with advancing age. The effect of diet is clearly shown, as calves fed more milk (AGG in Figure 2) maintained lower NDF digestion throughout the three digestibility periods. Also, calves fed functional fatty acids and nutrients (NeoTec5g®, Provimi North America, Brookville, OH, USA) as feed additives (MOD+ and AGG+ in Figure 2) had higher NDF digestion in periods 2 (42-46 d of age) and 3 (54 to 58 d of age). Previous studies (Guilloteau et al., 2009, 2010; Hill et al. 2007) have shown that feeding sodium butyrate (a component of NeoTec5g) improved fibre digestion in young calves.

Calves fed the moderate MR program (MOD in Figure 2) consumed more starter throughout the trial, which likely hastened rumen development and the ability of calves to digest NDF. In calves fed MOD, NDF digestion increased from approximately 15% at 19-23 d of age to approximately 35% by 51-56 d of age. Digestion of NDF in calves fed the higher level of MR (AGG) did not change markedly through the 56-d study and there were few differences with advancing age.




Figure 2. Change in total tract NDF digestibility in calves fed 0.66 kg of DM of a 27% CP, 17% fat MR powder daily fed for 49 d (MOD) without (-) or with (+) added NeoTec4 feed additive; or 0.66 kg of DM of a 27% CP, 17% fat MR powder fed for 4 d, then 0.96 kg of DM for 4 d, then 1.31 kg of DM fed for 34 d, then 0.66 kg of DM for 7 d (AGG). Effect of feeding level, NeoTec4 inclusion and age were significant ($P < 0.05$). Digestibility periods were 1 = 19-23 d; 2 = 40-44 d; and 3 = 52-56 d of the study. Calves were 2-3 d of age at initiation of the study. Adapted from Hill et al. (2016b).

In addition to age of calf, digestion of nutrients post-weaning is affected by ingredient source and form of calf starter. Digestion of DM, OM, and CP were higher in starters containing ground corn, whereas ADF and NDF digestion were greatest in starters containing soybean hulls (Table 4). Hill et al. (2016a) also reported that texturized calf starters containing whole corn and whole oats (51-54% starch and 13% NDF) had higher DM, OM, and CP digestibility than pelleted starters containing wheat middlings, soybean hulls, and dried distiller's grains (20% starch and 36% NDF; Table 5). On the other hand, pelleted, high-fibre starters had higher ADF, NDF, starch, and fat digestion. Gain of BW and hip width increased as OM digestibility increased in these trials.

Collectively, these data suggest that the availability of energy from starters is dependent on the type of carbohydrate, form of the starter (texturized vs. pelleted) and carbohydrate, age of the calf, and intake of liquid pre-weaning.

Table 4. Nutrient digestibility in calves 15-16 wk of age fed starters containing soybean hulls (S), wheat middlings (M) or corn (C). Contrast 1 = (S+M) vs. C; contrast 2 = S vs. M. Adapted from Hill et al., 2016a.

Digestibility, %	S	M	C	SE	Contrast 1	Contrast 2
DM	76.9	78.9	85.2	1.58	0.01	0.23
OM	77.5	79.6	85.8	1.56	0.01	0.21
ADF	65.5	53.5	55.4	3.48	0.20	0.01
NDF	70.7	56.1	66.2	3.13	0.34	0.01
Starch	97.6	98.9	97.0	0.57	0.13	0.15
CP	78.1	80.7	84.4	1.75	0.01	0.16
Sugar	94.2	95.6	94.2	1.79	0.63	0.47
Fat	84.1	86.3	89.6	2.61	0.08	0.42

Table 5. Nutrient digestibility in calves 15-16 wk of age fed high starch texturized (TX) or low starch pelleted (PL) starters containing low (MPL) or high MPH) amounts of metabolizable protein. No main effect of metabolizable protein was reported. P = probability of a main effect of starch level. Adapted from Hill et al., 2016a.

Digestibility, %	TX-MPL	TX-MPH	PL-MPL	PL-MPH	SEM	P
DM	84.3	84.7	79.7	78.8	0.51	0.001
OM	84.9	85.0	80.2	78.9	0.57	0.001
ADF	41.5	54.0	65.2	66.1	1.86	0.001
NDF	56.8	62.8	69.4	66.1	1.64	0.005
Starch	95.1	95.7	99.0	98.7	0.29	0.001
CP	84.9	84.6	79.5	78.6	0.54	0.001
Sugar	95.3	95.6	95.7	92.4	0.68	NS
Fat	86.3	82.7	88.3	87.8	0.78	0.08

Current nutrient models for calves and heifers (e.g., 2001 Dairy NRC) ignore the effects of previous nutrition and the extent of rumen development. The ME content of starters is a static calculation based on expected digestibility of nutrient fractions (NDF, non-fibre carbohydrate, protein, and fat). No provision is made for differing nutrient digestibilities with advancing age or intake. Conversely, other models for lactating cows utilize dynamic calculations of energy based on rates of ruminal digestion of each fraction (NFC, NDF, protein, fat) and rate of passage (Higgs et al., 2015). Intestinal

digestibility coefficients are then applied to the ruminally undegraded fraction to estimate total nutrient supply.

Using data from Chapman et al. (2016) and Hill et al. (2016b), we estimated ME concentrate of calf starter using the method outlined in the 2001 NRC Nutrient Requirements of Dairy Cattle (NRC, 2001) and calculated ME based on analyzed values using digestibility data from Table 3 and Figure 2. Results are in Table 6. The column labeled “NRC” contains calculated ME concentration in starter based on the 2001 NRC method, assuming digestibility values typical for adult ruminants. The column “Calculated” contains data using total tract digestibility measured in the studies by Chapman et al. (2016) and Hill et al. (2016b). We also used the 2001 Dairy NRC model to predict ME-allowable BW gain using the ME values calculated for calf starter using the NRC (NRC ME-g) or calculated values (Calc. ME-g) in Table 6.

Table 6. Estimated ME concentration (Mcal/kg of DM) in calf starters from Chapman et al. (2016) and Hill et al. (2016b) using methods of 2001 Dairy NRC (NRC) or calculated using total tract digestibilities reported in each experiment. ME-allowable BW gains were calculated using equations [2-4 a-e and 2-5 to 2-10] in 2001 Dairy NRC Requirements for Dairy Cattle (NRC, 2001) or using digestibility estimates from Table 3 and Figure 2, respectively. Digestibility estimates were made at 52-56 d.

Item	Starter ME, Mcal/kg			Predicted ME grain, kg/d		
	NRC	Calculated	%	NRC	Calc.	%
Chapman et al. 2016						
CON	2.81	2.59	92	0.77	0.67	87
MOD	2.81	2.56	91	0.93	0.82	88
AGG	2.84	2.30	81	0.94	0.70	74
Hill et al. 2016b						
MOD-	2.81	2.52	90	0.83	0.71	86
AGG-	2.89	2.45	85	0.61	0.45	74
MOD+	2.83	2.60	92	0.77	0.68	88
AGG+	2.87	2.50	87	0.70	0.55	79

Differences were significant for all measurements, but ME was markedly overestimated in calves fed higher levels of milk in both studies. Consequently, predicted ME-allowable gains using the calculated ME values for calf starter were lower compared to predicted gains using the ME values calculated with the NRC calculations.

The implications of calculation errors of ME content are clear, as calves fed high levels of milk pre-weaning will be ill prepared for weaning and unable to extract nutrients from calf starters efficiently. Consequently, growth of calves

will be compromised until sufficient maturation of the digestive tract and associated tissues allows the calf to fully utilize nutrients in the calf starter. The existing NRC model over-predicts ME supply from starters by 12 to 26% (Table 6).

These data also suggest that additional time may be needed for a weaning transition to ensure that calves fed high levels of milk will consume sufficient starter prior to weaning. In most of the studies cited in this review, liquid intake was reduced for 7-10 d prior to weaning. For calves fed 1 kg of powder or greater, this is probably insufficient time for adaptation.

■ Summary

The 2001 Dairy NRC represented an important improvement in our understanding of nutrient requirements for young calves and heifers. Further refinement of methods to estimate nutrient supply of young calves will improve our ability to calculate growth under a wide range of feeding and management conditions.

Feeding varying amounts of liquid from milk or MR has important implications to growth post-weaning. Increasing liquid consumption above approximately 650-700 g of solids per day will delay initiation of calf starter intake and will delay onset of rumen development. Digestion of all nutrients, particularly NDF, is essential to ensure that rumen development is adequate prior to weaning.

■ References

- Allen, M. S. 1997. Relationship between fermentation acid production in the rumen and the requirement for physically effective fiber. *J. Dairy Sci.* 80:1447–1462.
- Anderson K.L., T. G. Nagaraja, J. L. Morrill, T. B. Avery, S. J. Galitzer and J. E. Boyer. 1987. Ruminant microbial development in conventionally or early-weaned calves. *J. Anim. Sci.* 64:1215-1226.
- Baldwin VI, R.L., K.R. McLeod, J.L. Klotz, and R.N. Heitmann. 2004. Rumen development, intestinal growth and hepatic metabolism in the pre- and postweaning ruminant. 87(Suppl.):E55–E65.
- Chapman, C. E., P. S. Erickson, J. D. Quigley, T. M. Hill, H. G. Bateman II, F. X. Suarez-Mena, and R. L. Schlotterbeck. 2016. Effect of milk replacer program on calf performance and digestion of nutrients with age of the dairy calf. *J. Dairy Sci.* 99:2740–2747.
- Chapman, C. E., T. M. Hill, D. R. Elder, and P. S. Erickson. 2017. Nitrogen utilization, preweaning nutrient digestibility, and growth effects of Holstein dairy calves fed 2 amounts of a moderately high protein or conventional milk replacer. *J. Dairy Sci.* 100:279–292.

- Davis Rincker, L. E., M. J. VandeHaar, C. A. Wolf, J. S. Liesman, L. T. Chapin, and M. S. Weber Nielsen. 2011. Effect of intensified feeding of heifer calves on growth, pubertal age, calving age, milk yield, and economics. *J. Dairy Sci.* 94:3554–3567.
- Dennis, T. S., F. X. Suarez-Mena, T. M. Hill, J. D. Quigley, R. L. Schlotterbeck, and L. Hulbert. 2018. Effect of milk replacer feeding rate, age at weaning, and method of reducing milk replacer to weaning on digestion, performance, rumination, and activity in dairy calves to 4 months of age. *J. Dairy Sci.* 101:268–278.
- Diaz, M. C., M. E. Van Amburgh, J. M. Smith, J. M. Kelsey, and E. L. Hutten. 2001. Composition of growth of Holstein calves fed milk replacer from birth to 105-kilogram body weight. *J. Dairy Sci.* 84:830–842.
- Guilloteau, P., G. Savary, Y. Jaguelin-Peyrault, V. Romé, L. Le Normand, and R. Zabielski. 2010. Dietary sodium butyrate supplementation increases digestibility and pancreatic secretion in young milk-fed calves. *J. Dairy Sci.* 93:5842–5850.
- Guilloteau, P., R. Zabielski, and J. W. Blum. 2009. Gastrointestinal tract and digestion in the young ruminant: Ontogenesis, adaptations, consequences and manipulations. *J. Physiol. and Pharmacol.* 60(Suppl. 3):37-46.
- Higgs, R. J., L. E. Chase, D. A. Ross, and M. E. Van Amburgh. 2015. Updating the Cornell Net Carbohydrate and Protein System feed library and analyzing model sensitivity to feed inputs. *J. Dairy Sci.* 98:6340–6360.
- Hill, T. M., J. M. Aldrich, R. L. Schlotterbeck, and H. G. Bateman II. 2006a. Effects of feeding calves different rates and protein concentrations of twenty percent fat milk replacers on growth during the neonatal period. *Prof. Anim. Sci.* 22:252–260.
- Hill, T. M., J. M. Aldrich, R. L. Schlotterbeck, and H. G. Bateman II. 2006b. Effects of feeding rate and concentrations of protein and fat of milk replacers fed to neonatal calves. *Prof. Anim. Sci.* 22:374–381.
- Hill, T. M., J. M. Aldrich, R. L. Schlotterbeck, and H. G. Bateman II. 2006b. Effects of feeding rate and concentrations of protein and fat of milk replacers fed to neonatal calves. *Prof. Anim. Sci.* 22:374–381.
- Hill, T. M., J. M. Aldrich, R. L. Schlotterbeck, and H. G. Bateman II. 2007. Effects of changing the fatty acid composition of calf starters. *Prof. Anim. Sci.* 23:665-671.
- Hill, T. M., H. G. Bateman II, J. M. Aldrich, and R. L. Schlotterbeck. 2010. Effect of milk replacer program on digestion of nutrients in dairy calves. *J. Dairy Sci.* 93:1105–1115.
- Hill, T. M., H. G. Bateman II, J. M. Aldrich, J. D. Quigley, and R. L. Schlotterbeck. 2015. Inclusion of tallow and soybean oil to calf starters fed to dairy calves from birth to four months of age on calf performance and digestion. *J. Dairy Sci.* 98:4882–4888.

- Hill, T. M., J. D. Quigley, H. G. Bateman II, J. M. Aldrich, and R. L. Schlotterbeck. 2016a. Source of carbohydrate and metabolizable lysine and methionine in the diet of recently weaned dairy calves on digestion and growth. *J. Dairy Sci.* 99:2788–2796.
- Hill, T. M., J. D. Quigley, F. X. Suarez-Mena, H. G. Bateman II, and R. L. Schlotterbeck. 2016b. Effect of milk replacer feeding rate and functional fatty acids on dairy calf performance and digestion of nutrients. *J. Dairy Sci.* 99:6352–6361.
- Hoover, W. H. 1986. Chemical factors involved in ruminal fiber digestion. *J. Dairy Sci.* 69:2755–2766.
- Howarth, R.E., R.L. Baldwin, M. Ronning. 1968. Enzyme activities in liver, muscle, and adipose tissue of calves and steers. *J. Dairy Sci.* 51:1270–1274.
- Huber, J.T., N.L. Jacobson, and R.S. Allen. 1961. Digestive enzyme activities in the young calf. *J. Dairy Sci.* 44:1494–1501.
- Huber, J. T., 1969. Development of the digestive and metabolic apparatus of the calf. *J. Dairy Sci.* 52:1303–1315.
- Jasper, J. and D. M. Weary. 2002. Effects of ad libitum milk intake on dairy calves. *J. Dairy Sci.* 85:3054–3058.
- Leibholz, J. 1975. The development of ruminant digestion in the calf. I. The digestion of barley and soya bean meal. *Aust. J. Agric. Res.* 26:1081–1091.
- McCormick, R. J. and W. E. Stewart. 1966. Pancreatic secretion in the bovine calf. *J. Dairy Sci.* 50:568–571.
- Moallem, U., D. Werner, H. Lehrer, M. Zachut, L. Livshitz, S. Yakoby, and A. Shamay. 2010. Long-term effects of ad libitum whole milk prior to weaning and prepubertal protein supplementation on skeletal growth rate and first-lactation milk production. *J. Dairy Sci.* 93:2639–2650.
- Morrill, J. L., W. E. Stewart, R. J. McCormick and H. C. Fryer. 1970. Pancreatic amylase secretion by young calves. *J. Dairy Sci.* 53:72–78.
- National Research Council. 2001. *Nutrient Requirements of Dairy Cattle*. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- Quigley, III, J. D., C. G. Schwab, and W. E. Hylton. 1985. Development of rumen function in calves: nature of protein reaching the abomasum. *J. Dairy Sci.* 68:694–702.
- Roth, B. A., E. Hillmann, M. Stauffacher, and N. M. Keil. 2008. Improved weaning reduces cross-sucking and may improve weight gain in dairy calves. *Appl. Anim. Behav. Sci.* 111:251–261.
- Roth, B. A., N. M. Keil, L. Gyax, and E. Hillmann. 2009. Influence of weaning method on health status and rumen development in dairy calves. *J. Dairy Sci.* 92:645–656.
- Russell, J. B. and D. B. Wilson. 1996. Why are ruminal cellulolytic bacteria unable to digest cellulose at low pH? *J. Dairy Sci.* 79:1503–1509.
- Shriver B. J., W. H. Hoover, J. P. Sargent, R. J. Crawford Jr., and W. V. Thayne. 1986. Fermentation of a high concentrate diet as affected by ruminal pH and digesta flow. *J. Dairy Sci.* 69:413–419.

- Siddons, R. C. 1968. Carbohydrase activities in the bovine digestive tract. *Biochem. J.* 108:839-844.
- Strzetelski, J., B. Niwinska, J. Kowalczyk, and A. Jurkiewicz. 2001. Effect of milk replacer feeding frequency and level on concentrate intake and rearing performance of calves. *J. Anim. Feed Sci.* 10:413-420.
- Terré, M., M. Devant, and A. Bach. 2007. Effect of level of milk replacer fed to Holstein calves on performance during the preweaning period and starter digestibility at weaning. *Livestock Sci.* 110:82–88.
- USDA. 2016. Dairy 2014, "Dairy Cattle Management Practices in the United States, 2014". USDA-APHIS-VS-CEAH-NAHMS. Fort Collins, CO. #692.0216.



NOTES

