

Implications of Calf-To-Calf Transmission of *Mycobacterium Avium* Subspecies *Paratuberculosis* (MAP) for Johne's Disease Management

Caroline Corbett, Jeroen De Buck, Karin Orsel, Herman W. Barkema

Department of Production Animal Health, Faculty of Veterinary Medicine, University of Calgary. 3330 Hospital Dr NW Calgary AB T2N 4N1.
Email: cscorbet@ucalgary.ca

Johne's disease (JD) is a chronic, progressive, inflammatory disease caused by *Mycobacterium avium* subspecies *paratuberculosis* (MAP). In the subclinical phase, cows have decreased milk production, increasing their risk of being culled. JD is usually transmitted through ingestion of MAP-contaminated material, such as contaminated feed and milk. A common assumption is that only cows are infectious and only calves are susceptible; this has led to one of the reasons for separating calves from their dams and placing them in calf barns. The potential risk of calf-to-calf transmission is largely overlooked in current JD prevention and control programs. However, recent research has confirmed that calves can actively shed the bacterium. Therefore, placing calves together may not completely eliminate the spread of MAP. The aims of this study are to: 1) quantify the extent to which transmission of MAP occurs between penmates; and 2) determine onset and duration of MAP shedding after infection. We hypothesize that infectious calves will infect susceptible penmates and this will be detected in fecal and blood samples.

Materials and Methods: In our challenge experiment will involve 36 newborn calves obtained from MAP-negative heifers. Calves will be randomized into 8 groups of 4 calves. Each isolated housing unit will consist of 4 calves, 2 of which are inoculated, and 2 sentinel calves. During the 3 mo of group housing, blood, fecal and environmental samples will be collected 3 times a week, and extensive tissue sampling at the end of the trial to confirm and identify the extent of transmission.

Implications: The results obtained from this study will not only add to our understanding of MAP transmission, but also generate new knowledge regarding patterns of shedding and immune responses between penmates. With a better understanding of transmission in calves, more effective protocols can be implemented in order to mitigate or minimize the spread of MAP.